King's College
DOHA

Dear Parents,

Please note that it is mandatory for all children to have their vaccinations completed as per the Qatar protocol and to submit their completed vaccination booklet to the school they are joining in Qatar.

Any child who has not completed any mandatory vaccinations will need to arrange for this to be completed before starting school.

According to Hamad Medical Corporation's General Pediatrics Division , vaccines are administered to children in Qatar as below:

Schedule/Vaccines	At Birth	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	2 YEARS	4 - 6 YEARS	13 - 16 YEARS
BCG	BCG									
HEPATITIS B	HepatitisB									
HEXA		Hexa1	Hexa2							
PENTA				Penta						
TETRA						Tetra				
DTap									DTaPB	
Tdap										Tdap
ROTAVIRUS		Rota1	Rota2							
OPV			OPV1	OPV2			OPV(Booster1)		OPV(Booster2)	
PCV13		PCV1	PCV2	PCV3		PCV(Booster)				
MMR					MMR1		MMR2			
VARICELLA					Varicella1				Varicella2	
HEPATITIS A					HepA1		HepA2			
INFLUENZA				Influenza						
MENINGOCOCCAL ACYW135								Meningococcal		

- BCG and HBV0: given at birth.
- 6-in-1 vaccine known as Hexavalent, usually given at two months. It protects against diptheria, tetanus and pertussis (whooping cough), Hepatitis B, Polio and Haemophilus Influenzae type b disease (Hib).
- Pentavalent vaccine almost same like Hexavalent given at two, four and six months.
- MMR protects against measles, mumps and rubella (new recommendation - two times at 12 and 18 months).
- Hepatitis B series vaccine given at 24 months, 3-4 years, 4-6 years, 11-12 years.
- Chickenpox (varicella) vaccine given at 12 months and 3-4 years.
- Pneumococcal vaccine (given at two, four, six and 15 months) protects against pneumonia (chest infection), ear infections and meningitis (a brain infection).
- Rotavirus vaccine (given at two and four months) protects infants against rotavirus, the most common cause of serious diarrhoea in babies and young children.
- Meningococcal ACYW135 (given from two years) prevents four strains of the bacteria that cause Meningococcal disease: A, C, Y, and W-135.
- Influenza vaccine is recommended for children starting from two years.

King's College
DOHA

Immunisation schedule (based on the age of the child)

May we please ask you to check off the vaccines that have been administered to your child, attached a screenshot copy of their vaccination booklet page in the provided image box below, save the form, and email it to admissionsforms@kingscollegedoha.com

Pupil Name:

Vaccine	At Birth	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	2 Years	4-6 Years	13-16 Years
BCG	☐									
HEP B	☐									
HEXA		☐	☐							
PENTA				☐						
TETRA						☐				
Dtap									☐	
TDap										☐
ROTAV.		☐	☐							
OPV			☐	☐			☐		☐	
PCV13		☐	☐	☐		☐				
MMR					☐		☐			
VARICELLA					☐				☐	
HEP A					☐		☐			
INFLUENZA				☐						
M. ACYW135								☐		

We cannot stress enough the importance of ensuring that your child has followed and has been given the required vaccinations.

Without them they will not be eligible to attend school.

I hereby confirm and acknowledge that all information presented is true and up to date.

Signature:

Date:

Screenshot Upload Next Page